

# **Passive Acoustic Monitoring for Marine Mammals at Site A in the Cape Hatteras Study Area, May – December 2014**

A Summary of Work Performed by Amanda J. Debich, Simone Baumann-Pickering, Ana Širović, John A. Hildebrand, Ariel M. Brewer, Kait E. Frasier, Rohen T. Gresalfi, Sean T. Herbert, Sarah C. Johnson, Ally C. Rice, Leah M. Varga, and Sean M. Wiggins

Marine Physical Laboratory  
Scripps Institution of Oceanography  
University of California San Diego  
La Jolla, CA 92037

AND

A Summary of Work Performed by Lynne E.W. Hodge, Joy E. Stanistreet, and Andrew Read

Duke University Marine Laboratory  
135 Duke Marine Lab Road  
Beaufort, NC 28516

Prepared by Lynne Hodge, Joy Stanistreet, and Andrew Read

Duke University Marine Laboratory  
135 Duke Marine Lab Road  
Beaufort, NC 28516

Submitted to:  
The Department of the Navy  
Norfolk, VA

*Suggested Citation:*

Hodge, L., J. Stanistreet, and A. Read. 2016. Passive Acoustic Monitoring for Marine Mammals at Site A in the Cape Hatteras Survey Area, May – December 2014. Technical Report. Submitted to Naval Facilities Engineering Command (NAVFAC) Atlantic, Norfolk, Virginia, under Contract No. N62470-10-D-3011, Task Order CTO 0051, issued to HDR Inc., Norfolk, Virginia. Prepared 17 August 2016.

Individual technical reports of other HARP deployments are available at:  
<http://www.navy-marine-species-monitoring.us/reading-room/>

**This project is funded by US Fleet Forces Command and managed by Naval Facilities Engineering Command Atlantic as part of the US Navy's marine species monitoring program.**

## **Abstract**

A High-frequency Acoustic Recording Package (HARP; Wiggins and Hildebrand 2007) was deployed between May 2014 and April 2015 in the Cape Hatteras, NC, survey area at Site A in 850 m. This HARP sampled continuously at 200 kHz and recorded for 217 days between 9 May 2014 and 11 December 2014. The data were divided into three frequency bands (10 Hz – 1000 Hz, 500 Hz – 5000 Hz, and 1 kHz – 100 kHz) and scanned for marine mammal vocalizations using Long-Term Spectral Averages (LTSAs) and automated detectors. Vocalizations of blue whales, fin whales, minke whales, sei whales, North Atlantic right whales, humpback whales, *Kogia* spp., Risso's dolphins, sperm whales, Cuvier's beaked whales, Gervais' beaked whales, Blainville's beaked whales, and unidentified delphinids were detected in the data.

## Methods

The May – December 2014 Hatteras Site A HARP (Hatteras 04A) was deployed at  $35.34677^{\circ}$  N,  $74.84805^{\circ}$  W on 8 May 2014 (recording started on 9 May 2014) and recovered on 6 April 2015 (normal recording ended on 11 December 2014). The instrument location is shown in Figure 1. Bottom depth at the deployment site was approximately 850 m. A schematic diagram of the Hatteras 04A HARP is shown in Figure 2.




Figure 1. Location of Hatteras 04A HARP deployment in the Cape Hatteras survey area.

**May 2014 Hatteras Site A HARP  
as deployed**

Deployment: May 8, 2014  
Recovery: April 15, 2015  
Latitude: 35.34677 N  
Longitude: -74.84805 W  
Depth: 850m




Figure 2. Schematic diagram showing details of the Hatteras 04A HARP. Note that diagram is not drawn to scale.

Data were acquired continuously at a 200 kHz sampling rate during the Hatteras 04A deployment. This deployment provided a total of 5207 hours of data over the 217 days of recording.

The following methods are a summary of [Debich \*et al.\* \(2016\)](#). Members of the Scripps Whale Acoustics Lab manually scanned the data from the Hatteras 04A HARP deployment for marine mammal vocalizations and anthropogenic sounds (sonar, explosions, shipping, and airguns) using LTSAs. Automated computer algorithm detectors were also used to analyze the data. Personnel at Scripps Institution of Oceanography analyzed the data for all marine mammal vocalizations except for beaked whales. J.E. Stanistreet performed the analysis for beaked whales; these methods will be discussed later.

Prior to manual review of the data, LTSAs were made for three frequency bands: (1) 10 – 1000 Hz (with resolutions of 5 s in time and 1 Hz in frequency), (2) 10 – 5000 Hz (with resolutions of 5 s in time and 10 Hz in frequency), and (3) 1 – 100 kHz (with resolutions of 5 s in time and 100 Hz in frequency). For effective analysis of marine mammal and anthropogenic sounds, analysts scanned three frequency bands: (1) low-frequency, between 10-300 Hz, (2) mid-frequency, between 10-5000 Hz, and (3) high-frequency, between 1-100 kHz. Each band was analyzed for the sounds of an appropriate subset of species or sources. Blue, fin, sei, Bryde's, minke, and North Atlantic right whales as well as the 5-pulse signal were classified as low-frequency; humpback whales, shipping, explosions, airguns, underwater communications, low-frequency active sonar greater than 500 Hz, and mid-frequency active sonar were classified as mid-frequency; and the remaining odontocete and sonar sounds were considered high-frequency. Low-frequency sounds were analyzed in hourly bins; mid- and high-frequency vocalizations were analyzed in one-minute bins. Vocalizations were assigned to species when possible. For North Atlantic right whale calls, the data were only examined for up-calls. Information on the

detections of shipping, explosions, and underwater communications are not reported here but can be found in [Debich \*et al.\* \(2016\)](#).

Detections of most sounds were made by manually scanning LTSA. However, detectors were used for some calls, including fin whale 20-Hz calls, humpback whale calls, *Kogia* spp. clicks, and echolocation clicks from the family Delphinidae. Fin whale 20-Hz calls were detected using an energy detection method, which used a difference in acoustic energy between signal and noise, calculated from a 5 s LTSA with 1 Hz resolution. The frequency at 22 Hz was used as the signal frequency, while noise was calculated as the average energy between 10 and 34 Hz. The resulting ratio is termed the fin whale acoustic index and is reported as a daily average. All calculations were performed on a dB scale.

Humpback whale call detection effort was automated using an algorithm based on the generalized power law (Helble *et al.* 2012). After the generalized power-law algorithm was applied, a trained analyst verified the accuracy of the detected signals. No effort was made to separate song and non-song calls.

Three steps were involved in the classification of *Kogia* spp. clicks. First, clicks with energy between 70-100 Hz without energy in lower frequency bands were identified. Then, an expert system classified these clicks based on spectral characteristics and finally an analyst verified all echolocation click bouts manually as *Kogia* spp. clicks.

Echolocation clicks from the family Delphinidae were detected using a modified version of a Teager energy detector (Soldevilla *et al.* 2008, Roch *et al.* 2011). Events were reviewed manually to remove false detections. LTSAs were then manually examined to identify reoccurring echolocation click types. Clicks were manually classified into separate click types based on characteristics such as inter-click interval, spectral peaks/troughs, and peak frequency. Classification was carried out by comparison to species-specific spectral characteristics from HARP recordings in the Gulf of Mexico (Frasier 2015).

For analysis of beaked whale echolocation signals, an automated detection method customized for the Cape Hatteras HARP recordings was used. This method used the same initial automated detection steps described in detail in [Debich \*et al.\* \(2014\)](#) to find 75-second recording segments containing potential beaked whale frequency modulated pulses. A Teager Kaiser energy detector was used to find echolocation signals, and criteria based on peak and center frequency, duration, and sweep rate were used to discriminate between delphinid and beaked whale signals ([Debich \*et al.\* 2014](#)). Additional criteria based on the shape and duration of the signal envelope were then applied to reduce the high number of false detections of non-beaked whale clicks in the Cape Hatteras recordings. All detected signals with a signal envelope increasing after 20 sample points, and remaining above a 50 percent energy threshold for at least 19 sample points but no greater than 70 sample points were kept; signals not meeting these criteria were removed from the analysis. The remaining detections were grouped into detection events, with detections separated by no more than 5 minutes considered to be a single event. In a final computer-assisted manual classification step, each detected event was given a species label by a trained analyst, and any remaining false detections were rejected (as in Baumann-Pickering *et al.* 2013).



This method resulted in significantly more detections of beaked whales at Cape Hatteras than manual LTSA analysis for this site, due to the ability to detect faint, barely visible beaked whale clicks as well as beaked whale clicks mixed in with echolocation from other odontocete species.

### *Data Quality*

Normal recording ended on 11 December 2014. After that date, disk error issues caused skipping in the data. These disk error issues resulted in only four more files written on four different days between 26 December 2014 and 15 January 2015. These data were not analyzed.

### **Results**

Table 1 summarizes the detected and identified marine mammal vocalizations for the Hatteras 04A HARP deployment. Figures 3-17 show the daily occurrence patterns for the marine mammals detected in this dataset. Figure 18 shows the occurrence of mid-frequency active sonar. Figure 19 shows the occurrence of high-frequency active sonar. Figure 20 shows the occurrence of airguns. Underwater ambient noise during this deployment is shown in Figure 21.

Mysticete detections included blue whales, fin whales, minke whales, sei whales, North Atlantic right whales, and humpback whales. Blue whale calls were detected only in October (Figure 3). Fin whale 20-Hz pulses (as measured by the acoustic index) were detected throughout the deployment, with peaks in calling in November and December (Figure 4). Minke whale pulse trains were first detected starting in October and continued through the end of the recording

period (Figure 5). Sei whale downsweeps were detected only in November (Figure 6). North Atlantic right whale up-calls were detected on only one day, 14 November 2014 (Figure 7). The timing coincides with the migration of this species to the breeding grounds. Humpback whale calls were detected on five days between May and November during this deployment (Figure 8).

Detected odontocete vocalizations included clicks and whistles (Figures 9-17). Many of these detections were assigned to the unidentified odontocete category, with whistles divided into two categories based on frequency (Figures 9-10) and with unidentified clicks being divided into four main groups based on spectral patterns (Figure 11). These vocalizations were present nearly continuously throughout the deployment. For more details on each of the four groups of clicks and which species may have produced them, see [Debich \*et al.\* \(2016\)](#). Clicks produced by *Kogia* spp. were also detected throughout the deployment (Figure 12). Risso's dolphin clicks were detected on a total of five days – three days in May and two days in September (Figure 13). Sperm whales were detected throughout the deployment during both day and night, with peaks in click detections in the summer months (Figure 14). There were also several click detections that were assigned to beaked whales. Cuvier's beaked whale clicks occurred regularly throughout the deployment, with detections distributed fairly uniformly across both seasonal and diel time scales (Figure 15). Gervais' beaked whale clicks occurred less frequently, with most detections between the end of June and July and between the end of October and the end of the recording period (mid-December) (Figure 16). Finally, Blainville's beaked whale clicks were detected on only two days in early December (Figure 17).

Table 1. Summary of detections of marine mammal vocalizations at the Cape Hatteras Survey Area Site A for May – December 2014 (Hatteras 04A). Fin whale 20-Hz pulses are not included as they were reported as an acoustic index and not logged with a start and end time to individual detection events.

| <b>Species</b>                          | <b>Call type</b>                           | <b>Total duration of vocalizations (hours)</b> | <b>Percent of recording duration</b> | <b>Days with vocalizations</b> | <b>Percent of recording days</b> |
|---|--|--|--------------------------------------|--------------------------------|----------------------------------|
| Blue whale <sup>a</sup>                 | A and B calls                              | 22   | 0.42                                 | 4                              | 1.84                             |
| Minke whale <sup>a</sup>                | pulse train (slow-down, speed-up, regular) | 237  | 4.55                                 | 35                             | 16.13                            |
| Sei whale <sup>a</sup>                  | downsweep                                  | 15   | 0.29                                 | 7                              | 3.23                             |
| North Atlantic right whale <sup>a</sup> | up-call                                    | 2  | 0.04                                 | 1                              | 0.46                             |
| Humpback whale                          | variable                                   | 0.45   | 0.009                                | 5                              | 2.30                             |
| Unidentified odontocete                 | whistles                                   | 955.88   | 18.36                                | 196                            | 90.32                            |
| Unidentified odontocete                 | clicks                                     | 1122.60  | 21.56                                | 210                            | 96.77                            |
| <i>Kogia</i> spp.                       | clicks                                     | 1.87   | 0.04                                 | 39                             | 17.97                            |
| Risso's dolphin                         | clicks                                     | 5.87   | 0.11                                 | 5                              | 2.30                             |
| Sperm whale                             | clicks                                     | 571.37   | 10.97                                | 97                             | 44.70                            |
| Cuvier's beaked whale                   | clicks                                     | 231.22   | 4.44                                 | 210                            | 96.77                            |
| Gervais' beaked whale                   | clicks                                     | 29.15  | 0.56                                 | 87                             | 40.09                            |
| Blainville's beaked whale               | clicks                                     | 0.08   | 0.002                                | 2                              | 0.92                             |

<sup>a</sup>Analyzed in hourly bins versus one-minute bins.




Figure 3. Blue whale call detections (black bars) in hourly bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 4. Weekly value of fin whale 20-Hz call acoustic index for the Hatteras 04A deployment.




Figure 5. Minke whale pulse train detections (black bars) in hourly bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 6. Sei whale downsweep detections (black bars) in hourly bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 7. North Atlantic right whale up-call detections (black bars) in hourly bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 8. Humpback whale call detections (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 9. Unidentified odontocete whistle detections that were less than 5 kHz (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 10. Unidentified odontocete whistle detections that were greater than 5 kHz (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 11. Unidentified odontocete click detections (different colored horizontal bars represent the different groups clicks were divided into for this report, with those in yellow not assigned a category) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 12. *Kogia* spp. click detections (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).






Figure 13. Risso's dolphin click detections (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 14. Sperm whale click detections (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 15. Cuvier's beaked whale click detections (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 16. Gervais' beaked whale click detections (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 17. Blainville's beaked whale click detections (black bars) in one-minute bins for the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 18. Mid-frequency active sonar (black bars) detected during the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 19. High-frequency active sonar (black bars) detected during the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 20. Airgun detections (black bars) within the Hatteras 04A deployment. Dark gray shading indicates periods of darkness, determined from the U.S. Naval Observatory (<http://aa.usno.navy.mil>).




Figure 21. Monthly averages of ambient noise at Cape Hatteras, NC, Site A for May – December 2014. Months with an asterisk (\*) are partial recording periods. Figure from Debich *et al.* (2016).

## References

- Baumann-Pickering, S., M.A. McDonald, A.E. Simonis, A. Solsona Berga, K.P.B. Merkins, E.M. Oleson, M.A. Roch, S.M. Wiggins, S. Rankin, T.M. Yack, and J.A. Hildebrand. 2013. Species-specific beaked whale echolocation signals. *Journal of the Acoustical Society of America* **134**: 2293-2301.
- Debich, A.J., S. Baumann-Pickering, A. Širović, J.A. Hildebrand, A.M. Brewer, K.E. Frasier, R.T. Gresalfi, S.T. Herbert, S.C. Johnson, A.C. Rice, L.M. Varga, S.M. Wiggins, L.E.W. Hodge, J.E. Stanistreet, and A.J. Read. 2016. [Passive acoustic monitoring for marine mammals in the Virginia Capes Range Complex October 2012 – April 2015. Final Report, Marine Physical Laboratory Technical Memorandum #559](#). August 2016. Submitted to Naval Facilities Engineering Command (NAVFAC) Atlantic, Norfolk, Virginia, under Contract No. N62470-10-D-3011 – Task Order Number 051 – issued to HDR, Inc.
- Debich A.J., S. Baumann-Pickering, A. Širović, J.S. Buccowich, Z.E. Gentes, R.S. Gottlieb, S.C. Johnson, S.M. Kerosky, L.K. Roche, B. Thayre, J.T. Trickey, S.M. Wiggins, J.A. Hildebrand, L.E.W. Hodge, and A.J. Read. 2014. [Passive Acoustic Monitoring for Marine Mammals in the Cherry Point OPAREA 2011-2012. Final Report, Marine Physical Laboratory Technical Memorandum #545](#). Submitted to Naval Facilities Engineering Command (NAVFAC) Atlantic, Norfolk, Virginia, under Contract No. N62470-10D-3011 issued to HDR, Inc.
- Frasier, K.E. 2015. Density estimation of delphinids using passive acoustics: A case study in the Gulf of Mexico. Doctoral dissertation, University of California San Diego, Scripps Institution of Oceanography, La Jolla, CA, USA. 321 pp.
- Roch, M.A., H. Klinch, S. Baumann-Pickering, D.K. Mellinger, S. Qui, M.S. Soldevilla, and J.A. Hildebrand. 2011. Classification of echolocation clicks from odontocetes in the Southern California Bight. *Journal of the Acoustical Society of America* **129**: 467-475.
- Wiggins, S.M. and J.A. Hildebrand. 2007. High-frequency Acoustic Recording Package (HARP) for broad-band, long-term marine mammal monitoring. In: *International Symposium on Underwater Technology 2007 and International Workshop on Scientific Use of Submarine Cables & Related Technologies 2007*: 551-557. Tokyo, Japan: Institute of Electrical and Electronics Engineers.